

LEVEL ^H

(R)

OFFICE OF NAVAL RESEARCH

Contract N00014-79-C-0647

Technical Report No. 2

AD A 096396

TWO PHOTON SPECTROSCOPY OF UF_6 IN THE NEAR ULTRAVIOLET


by

E. R. BERNSTEIN AND P. M. KENNEDY

Prepared for Publication in
The Journal of Chemical Physics

Department of Chemistry
Colorado State University
Fort Collins, Colorado 80523

29 August 1980



Reproduction in whole or in part is permitted for
any purpose of the United States Government.

Approved for Public Release; Distribution Unlimited.

DBC FILE COPY

81 3 16 126

UNCLASSIFIED


SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 2	2. GOVT ACCESSION NO. AD-A096896	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) (6) TWO PHOTON SPECTROSCOPY of UF_6 IN THE NEAR ULTRAVIOLET.	5. TYPE OF REPORT & PERIOD COVERED (9) TECHNICAL REPORT.	
7. AUTHOR(s) (10) E.R./Bernstein and P.M./Kennedy	8. CONTRACT OR GRANT NUMBER(s) (13) N00014-79-C-0647	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Colorado State University Fort Collins, Colorado 80523 (12) 12/	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS (11) 18 Mar 81	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, Virginia 22217 (14) TR-2	12. REPORT DATE March 10, 1981	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES 11	
	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release, Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Spinorbit coupling; hexahalides; dye-laser; super-quadronic multiphoton absorption; emission signals; two photon spectroscopy; photon acoustic spectroscopy; fluorescent spectroscopy.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) → The two photon excited fluorescence excitation spectrum of UF_6 has been observed in the region 410 to 315 nm. The spectrum is virtually indistinguishable from the one photon absorption in this region. No vibronic structure was observed in absorption or dispersed emission. These data indicate a high density of u and g states for UF_6 in this energy range and facile photochemical decomposition. Emission intensity is found to be proportional to laser power to the 3.0 to 3.6 power.		

I. Introduction

Uranium hexafluoride has been the subject of intense spectroscopic investigation for the past 40 years.¹⁻⁹ The one-photon gas phase absorption spectrum exhibits only broad charge transfer bands with maxima at 214, 260 and 368 nm. To a first approximation, these bands may be considered as resulting from an electron being promoted from a fluorine 2p orbital to a uranium 5f orbital. Lewis and coworkers⁴ have published a more detailed band assignment which takes into account the strong spin-orbit coupling in UF_6 . In their scheme, the uppermost filled molecular orbital (primarily a ligand t_{1u} orbital) transforms as T_{2u} , and the five lowest (primarily metal) unfilled orbitals transform as T_{7u} , T_{8u} , T_{7u} , T_{8u} , and T_{6u} in order of increasing energy. Thus, the 260 and 368 nm bands are assigned to electric dipole forbidden, magnetic dipole allowed $u \rightarrow u$ transitions, while the first electric dipole allowed transition occurs from the molecular orbital immediately below the highest filled MO (a T_{8g} orbital) and is identified with the 214 nm band. It would therefore appear profitable to study multiphoton, and in particular two-photon, spectroscopy of UF_6 since the selection rules are complementary to the selection rules for one-photon spectroscopy.

Low temperature solid state UF_6 emission and excitation spectra exhibit detailed vibronic structure.^{5,9} The one-photon gas phase absorption spectrum of MoF_6 , a molecule similar to UF_6 , has been reported to show a progression in ν_1 in the ultraviolet.^{10,3} Thus, the possibility of observing two-photon vibronic structure provides a further motivation for two-photon absorption experiments with UF_6 .



A third impetus for these investigations is provided by Lewis's² molecular orbital diagram which places the two highest occupied molecular orbitals in UF_6 in the reverse order from that observed in most other hexahalides. Thus, t_{1g}^{π} is predicted to be below t_{1u}^{σ} in UF_6 whereas t_{1u}^{σ} is the more stable in most other hexahalides. This point is discussed in a review by Jorgensen¹¹. Two-photon spectroscopy might reasonably be expected to provide additional information regarding this reversal.

Finally, elucidation of the low lying electronic and vibronic states of UF_6 would be an important step towards an understanding of its photo-physical and photochemical behavior.

II. Experimental

The experimental apparatus used in this investigation has been described previously^{12,13}. Briefly, the purified (triply distilled in vacuo) UF_6 is contained in a stainless steel can which is attached to the optical cell with stainless steel tubing. The cell is constructed of stainless steel and quartz windows with indium seals. A beam from a Nd/YAG-pumped dye laser (Quanta Ray) is focused into the cell and fluorescence is observed at 90° to the beam direction via an RCA 8850 photomultiplier tube. The signal from the PM tube is fed into channel A of a PAR Boxcar Integrator. The beam, after passing through the optical cell, impinges upon a power meter, the signal from which is fed into channel B of the boxcar. Spectra are recorded as A/B^2 in an attempt to normalize intensity to dye output and minimize pulse-to-pulse laser variations. A CuSO_4 filter is placed between the cell and the PM tube to remove scattered laser light, and suitable sharp-cut filters are placed on the cell to remove unwanted dye-laser harmonics and incoherent fluorescence. To obtain sufficient power in the low energy region of the spectrum, a Raman-shifter is employed with some of the dyes to shift the wavelength by 4155 cm^{-1} . The Raman shifter contains hydrogen at a pressure of 100 psi. The following dyes are used to scan the wavelength ranges indicated:

DCM	628-670 nm
Carbazine 720	684-720
Rhodamine 590 (Raman-shifted)	716-750
Rhodamine 610 (Raman-shifted)	746-774
Kiton Red (Raman-shifted)	760-784
Rhodamine 640 (Raman-shifted)	794-816
DCM (Raman-shifted)	815-910

In the DCM and Carbazine 720 dye regions, absorption spectra were also obtained photoacoustically as described in reference 12. Essentially the same two-photon spectra are obtained by both techniques.

In addition to the absorption spectra, an attempt was made to disperse the UF_6 emission upon excitation at two separate wavelengths: 640 nm (DCM dye - 260 nm band) and 775 nm (Kiton Red dye - 368 nm band). Excitation at 640 nm yields a weak continuous emission over the wavelength range 300 to 450 nm. No structure whatsoever was observed. Excitation at 775 nm yielded an emission signal too weak to be dispersed.

III. Results and Discussion

If it is assumed that two photons are absorbed simultaneously, the wavelength range covered corresponds to a range which includes the low energy side of the 260 nm band to the low energy side of the 368 nm band of the one-photon spectrum (342-410 nm). In the one-photon spectrum, a minimum is observed in this region at 340 nm, with strong absorption to high energy and weak absorption to the low energy side of this minimum. No absorption is observed above 410 nm^{3,4}. The multiphoton (fluorescence) excitation spectrum exhibits the same overall shape as the one-photon spectrum in the wavelength range studied with the minimum shifting to 354 nm. Strong absorption is observed to high energy and weak absorption to the low energy side of this minimum. No absorption is observed above 410 nm. Similar results obtain for the photoacoustic spectra.

In order to determine the number of photons simultaneously absorbed, a power dependence study was performed on several of the dyes (R610, R590 and C720) by plotting fluorescence intensity vs. laser power on a log-log graph. The slopes of the lines obtained were generally between 3.0 and 3.6 suggesting the absorption of 3 or 4 photons. This is believed to be the result of two effects: (1) Real, super-quadratic multiphoton absorption and (2) the build-up of a UF_5 or $(UF_5)_n$ "snow" within the cell which strongly scattered the incoming beam and lead to a spuriously high "fluorescence" intensity.¹⁴ UF_6 is known to photolyze rapidly to UF_5 which then polymerizes.¹⁵ The sensitivity of the PM tube was sufficiently high that even a few photons passing through the $CuSO_4$ filter would yield higher than normal intensities. As more experiments were performed between cell cleanings, the number of photons apparently absorbed became higher, indicating a build-up of material within the cell.

The above mentioned difficulties notwithstanding, we believe that both photoacoustic and emission signals originate with resonant two-photon absorption in the region 340-410 nm for the following two reasons: (1) no absorption is observed on the low energy side of 410 nm even though there are states which would be resonant for greater-than-two photon absorption; and (2) the interpretation of the spectra of other molecules (pyrazene¹², triazine¹³ and osmium tetroxide¹⁶) in the same experimental apparatus using similar laser power levels is only possible by assuming a two-photon resonant absorption even though emission may arise from higher states and/or photochemically-generated species.

Polarization studies were performed using both photo electric and photo acoustic detection. However, little significant difference could be found between the two spectra. This is to be expected considering there is so little difference between one and two photon spectra and that there are no pronounced features in either spectra.

IV. Conclusions

Two-photon absorption spectroscopy of gas phase UF_6 yields a spectrum which, in its main features, is indistinguishable from the one-photon absorption spectrum. No vibronic structure is observed. Two-photon excited emission is weak and continuous for the 260 nm band and about a factor of 10 weaker for the 360 nm band. Again, no vibronic structure is observed. These results are indicative of the difficulty in performing experiments on a molecule which rapidly and efficiently photolyzes. However, the similarity of the one and two-photon spectra does indicate that the density of excited states accessible to UF_6 must be very high in this energy range.

Investigations of the solid state of UF_6 appear much more promising in this regard. Well-resolved emission and excitation spectra have been obtained in matrix-isolated and pure and mixed crystal UF_6 . The solid state and low temperatures employed appear to suppress the fast photochemistry to a significant extent.

References

1. G. H. Dieke and A.B.F. Duncan, "Spectroscopic Properties of Uranium Compounds" (McGraw-Hill, New York, 1949), Chap. 8.
2. G. D. Sheremet'ev, Opt. Spektrosk., 2, 99 (1957) and 1, 181 (1956).
3. R. McDiarmid, J. Chem. Phys., 65, 168 (1976).
4. W. B. Lewis, L. B. Asprey, L. H. Jones, R. S. McDowell, S. W. Rabideau, A. H. Zeltmann, J. Chem. Phys., 65, 2707 (1976).
5. J. C. Miller, S. W. Allison, L. Andrews, J. Chem. Phys., 70, 3524 (1979).
6. R. C. Oldenberg, W. W. Rice, F. B. Wampler, J. Chem. Phys., 69, 2181 (1978).
7. W. W. Rice, F. B. Wampler, R. C. Oldenberg, W. B. Lewis, J. J. Tiee, R. T. Pack, J. Chem. Phys., 72, 2948 (1980).
8. W. B. Lewis, A. H. Zeltmann, J. Photochem., 12, 51 (1980).
9. E. R. Bernstein, G. R. Meredith, J. D. Webb, J. Chem. Phys., 68, 4066 (1978).
10. K. N. Tanner, A.B.F. Duncan, J. Am. Chem. Soc., 73, 1164 (1951).
11. C. K. Jorgensen, Prog. Inorg. Chem., 12, 101 (1970).
12. J. D. Webb, K. M. Swift, E. R. Bernstein, J. Mol. Struct., 61, 285 (1980).
13. J. D. Webb, K. M. Swift, E. R. Bernstein, J. Chem. Phys., 73, (1980).
14. Similar results have recently been reported: E. Borsella, F. Catoni, G. Freddi, J. Chem. Phys. 73, 316 (1980).
15. W. B. Lewis and A. H. Zeltmann, J. Photochem. 12, 51 (1980).
16. E. R. Bernstein, K. M. Swift, unpublished results.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Area Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
ONR Western Regional Office 1020 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
ONR Eastern/Central Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 3100 (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Dr. John Doyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, 051A

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. M. A. El-Sayed Department of Chemistry University of California, Los Angeles Los Angeles, California 90024	1	Dr. M. Rauhut Chemical Research Division American Cyanamid Company Bound Brook, New Jersey 08805	1
Dr. E. R. Bernstein Department of Chemistry Colorado State University Fort Collins, Colorado 80521	1	Dr. J. I. Zink Department of Chemistry University of California, Los Angeles Los Angeles, California 90024	1
Dr. C. A. Heller Naval Weapons Center Code 6059 China Lake, California 93555	1	Dr. D. Haarer IBM San Jose Research Center 5600 Cottle Road San Jose, California 95143	1
Dr. J. R. MacDonald Chemistry Division Naval Research Laboratory Code 6110 Washington, D.C. 20375	1	Dr. John Cooper Code 6130 Naval Research Laboratory Washington, D.C. 20375	1
Dr. G. B. Schuster Chemistry Department University of Illinois Urbana, Illinois 61801	1	Dr. William M. Jackson Department of Chemistry Howard University Washington, DC 20059	1
Dr. A. Adamson Department of Chemistry University of Southern California Los Angeles, California 90007	1	Dr. George E. Walraffen Department of Chemistry Howard University Washington, DC 20059	1
Dr. M. S. Wrighton Department of Chemistry Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Dr. Rudolph J. Marcus Office of Naval Research Scientific Liaison Group American Embassy APO San Francisco 96503	1
		Mr. James Kelley DTNSRDC Code 2803 Annapolis, Maryland 21402	1